

## Birmingham, Alabama — 1999 (N=300)

Figure A. Age of GISP participants, in years, 1999




Figure B. Race/ethnicity of GISP participants, 1999




Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988–1999






Figure D. Reason for visit among GISP participants, 1999



## Birmingham, Alabama — 1999 (N=300)

**Figure E. Previous episode of gonorrhea among GISP participants, 1991–1999**




\*Data first collected in 1991.


\*\*Data first collected in 1992.

Note: Data points not shown when >30% data missing.


**Figure F. Drugs used to treat gonorrhea among GISP participants, 1999**



**Figure G. Drugs used to treat chlamydia trachomatis infection among GISP participants, 1999**



**Figure H. Resistance to penicillin and tetracycline among GISP isolates, 1999**




## Birmingham, Alabama — 1999 (N=300)


**Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988–1999**

No isolates with decreased susceptibility to ceftriaxone have been identified at this clinic.

**Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992–1999**



**Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990–1999**



Note: Susceptibility to ciprofloxacin first measured in 1990.

**Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992–1999**

No isolates with decreased susceptibility to azithromycin have been identified at this clinic.